

On the Spectrum of $D = 2$ Supersymmetric Yang-Mills Quantum Mechanics

Yoji Michishita *

*Department of Physics, Faculty of Education, Kagoshima University
Kagoshima, 890-0065, Japan*

March, 2011

Abstract

We investigate the structure of the spectrum of states in $D = 2$ $SU(N)$ supersymmetric Yang-Mills matrix quantum mechanics, which is a simplified model of Matrix theory. We compute the thermal partition function of this system and give evidence for the correctness of naively conjectured structure of the spectrum. It also suggests that Claudson-Halpern-Samuel solution is the unique eigenfunction of simultaneously diagonalizable hermitian operators, and we show that it is true in $N = 3$ and $N = 4$ cases.

*michishita@edu.kagoshima-u.ac.jp

1 Introduction

Matrix theory is expected to be a correct description of M-theory, and is given as an $U(N)$ matrix quantum mechanics. The structure of the spectrum of states of this theory is complicated, and we know little about it except that the spectrum is continuous with some discrete poles.

Instead, in this paper we investigate $D = 2$ $SU(N)$ supersymmetric Yang-Mills matrix quantum mechanics, whose action is given by the dimensional reduction of $D = 2 = 1 + 1$ super Yang-Mills theory to $0 + 1$ dimension. As a simplified model of Matrix theory, which is from $D = 10$ super Yang-Mills theory, we expect that it gives some hint to the spectrum of Matrix theory. This quantum mechanics is almost free, but Gauss law associated with the gauge fixing procedure makes it nontrivial.

The action of $D = 2$ matrix quantum mechanics is given by

$$S = \frac{1}{g^2} \int dt \operatorname{tr} \left[\frac{1}{2} DXDX + ig^2 \theta^\dagger D\theta - g^4 \theta^\dagger [X, \theta] \right], \quad (1.1)$$

where g is the coupling constant, $DX = \partial_t X - i[A_0, X]$, and X, A_0, θ are matrices in $su(N)$ Lie algebra. θ is also a complex Grassmann odd operator. This action is invariant under the gauge transformation, and the following supersymmetry transformation:

$$\delta X = i(\epsilon^\dagger \theta + \epsilon \theta^\dagger), \quad \delta \theta = -\Pi \epsilon, \quad \delta A_0 = g^2 \delta X. \quad (1.2)$$

Quantization of this system gives the following commutation relations:

$$[X^a, \Pi^b] = i\delta_{ab}, \quad \{\theta^a, (\theta^b)^\dagger\} = \delta_{ab}, \quad (1.3)$$

where $\Pi = \frac{1}{g^2} DX$ and a is an index of $SU(N)$ adjoint representation. Physical states must be gauge invariant: $G^a |\text{phys}\rangle = 0$, where G is the generator of $SU(N)$:

$$G = i[X, \Pi] + \{\theta^\dagger, \theta\}. \quad (1.4)$$

Hamiltonian H and supercharge Q are

$$H = g^2 \operatorname{tr} \left[\frac{1}{2} \Pi^2 \right] + (\text{terms proportional to 1st class constraints}), \quad (1.5)$$

$$Q = -\operatorname{tr} [\Pi \theta]. \quad (1.6)$$

On physical states H acts as free Hamiltonian $h = g^2 \operatorname{tr} [\frac{1}{2} \Pi^2]$. However, this system is not trivial, because of the physical condition $G |\text{phys}\rangle = 0$. Commutation relations of H and Q

are usual ones, up to operators which vanish on physical states:

$$\{Q^\dagger, Q\} = \frac{2}{g^2}h + (\text{terms proportional to 1st class constraints}), \quad (1.7)$$

$$\{Q, H\} = (\text{terms proportional to 1st class constraints}), \quad (1.8)$$

$$\{Q^\dagger, H\} = (\text{terms proportional to 1st class constraints}). \quad (1.9)$$

The fermion part of the states is constructed by acting $(\theta^a)^\dagger$ on a vacuum $|0_F\rangle$ satisfying $\theta^a|0_F\rangle = 0$. Then the total states take the following form:

$$(\theta^{a_1})^\dagger \dots (\theta^{a_n})^\dagger |0_F\rangle \int \prod_a dX^a \left[\prod_a |X^a\rangle \right] \psi(X). \quad (1.10)$$

A gauge invariant solution to Schrödinger equation in the sector of zero fermion number has been found in [1, 2]: When X is diagonalized by the gauge transformation:

$$X = U \begin{pmatrix} x_1 & & & \\ & x_2 & & \\ & & \ddots & \\ & & & x_N \end{pmatrix} U^{-1}, \quad (1.11)$$

then Claudson-Halpern-Samuel (CHS) solution, which has N continuous parameters k_i , is given by

$$\psi(X) = \frac{1}{\mathcal{M}} \sum_{\sigma} \text{sgn}(\sigma) \exp \left[i \sum_{i=1}^N k_{\sigma(i)} x_i \right], \quad (1.12)$$

where $\mathcal{M} = \prod_{i < j} (x_i - x_j)$ and $\sum_{\sigma}$ indicates summation over all the permutations. This expression gives a solution in the case of $U(N)$ gauge group. To remove center of mass $U(1)$ part one just have to impose the condition $\sum_i k_i = 0$. Then $\psi(X)$ depends on only traceless part $w_i \equiv x_i - \frac{1}{N} \sum_j x_j$. In $SU(2)$ case this is the only energy eigenfunction with zero fermion number[1, 2]. In the case of higher groups there are more solutions [3, 4].

Naively we can expect that the spectrum can be constructed by acting gauge invariant operators made of Π and $\theta^\dagger$ on states in the sector of zero fermion number[2]. Indeed in $N = 2$ case it has been proven that the whole spectrum is constructed in this way on CHS solution[1, 2]. To investigate the case of general N , in section 2 we compute the thermal partition function of this system, and give an interpretation of the result. The form of the partition function gives support to the naive expectation, and we give a conjecture that CHS solution is the unique eigenfunction of simultaneously diagonalizable operators $\text{tr} [\Pi^n]$ ($n = 2, 3, \dots, N$). In section 3, we prove this conjecture in $N = 3$ and $N = 4$ cases. Section 4 contains some discussion, and Appendix contains necessary information on $SU(N)$ group theory.

2 Thermal partition function and spectrum of $D = 2$ matrix quantum mechanics

In this section we compute the thermal partition function of $D = 2$ quantum mechanics $Z = \text{Tr}_{\text{phys}} [e^{-\beta H}]$, where the trace Tr_{phys} is taken over the physical states. Then from it we will obtain insight to the structure of the spectrum of this system. First we compute Z in the operator formalism. Z can be rewritten in terms of a trace without the condition of gauge invariance, by inserting $\text{SU}(N)$ group element $e^{i\lambda^a G^a}$:

$$Z = \int d\lambda \text{Tr} [e^{-\beta h + i\lambda^a G^a}], \quad (2.1)$$

where $\int d\lambda$ indicates integral over $\text{SU}(N)$ (See Appendix for notation about $\text{SU}(N)$ group theory used here and the following), and works as a projector onto the gauge invariant states. We can set $\lambda^{(ij)} = 0$ without loss of generality.

Since in $-\beta h + i\lambda^a G^a$ boson part and fermion part are decoupled, Z is decomposed into corresponding parts Z_B and Z_F :

$$Z = \int d\lambda Z_B(\lambda) Z_F(\lambda), \quad Z_B(\lambda) = \text{Tr} [e^{-\beta h + i\lambda^a G_B^a}], \quad Z_F(\lambda) = \text{Tr} [e^{i\lambda^a G_F^a}], \quad (2.2)$$

where

$$G_B^a = f_{bac} X^b \Pi^c, \quad G_F^a = -i f_{bac} (\theta^b)^\dagger \theta^c. \quad (2.3)$$

Z_F can be computed by using the knowledge of group theory: $e^{i\lambda^a G_F^a}$ is just the gauge group element for fermions:

$$e^{i\lambda^a G_F^a} (\theta_b)^\dagger e^{-i\lambda^a G_F^a} = [\text{Ad}(\lambda)]_{bc} (\theta_c)^\dagger, \quad (2.4)$$

and $(\theta^{a_1})^\dagger \dots (\theta^{a_n})^\dagger |0_F\rangle$ forms an antisymmetric tensor product representation of adjoint representation. Therefore Z_F is given by the sum of group theory characters of those representations:

$$Z_F(\lambda) = \sum_{n=0}^{N^2-1} \chi(\text{Alt}_n(\text{adj.})) = 2^{N-1} \prod_{\substack{i,j=1 \\ i \neq j}}^N (1 + z_i/z_j), \quad (2.5)$$

where we used (A.23). If we introduce an additional parameter q for counting fermion number F : $Z_F(\lambda, q) \equiv \text{Tr} [q^F e^{i\lambda^a G_F^a}]$, then similarly,

$$Z_F(\lambda, q) = \sum_{n=0}^{N^2-1} q^n \chi(\text{Alt}_n(\text{adj.})) = (1+q)^{N-1} \prod_{\substack{i,j=1 \\ i \neq j}}^N (1 + q z_i/z_j). \quad (2.6)$$

Next let us compute $Z_B(\lambda)$. By introducing eigenstates of Π : $\Pi^a |p_b\rangle = p^a |p_b\rangle$,

$$\begin{aligned} Z_B(\lambda) &= \left[\prod_a \int_{-\infty}^{\infty} \frac{dp_a}{2\pi} \right] \langle p_b | e^{-\beta h} e^{i\lambda^c G_B^c} | p_b \rangle \\ &= \left[\prod_a \int_{-\infty}^{\infty} \frac{dp_a}{2\pi} \right] e^{-\frac{\beta g^2}{2} p_d^2} \langle p_b | e^{i\lambda^c G_B^c} | p_b \rangle. \end{aligned} \quad (2.7)$$

Noting that $e^{i\lambda^c G_B^c}$ generates gauge group elements,

$$\langle p_b | e^{i\lambda^c G_B^c} | p_b \rangle = \langle p_b | \text{Ad}(\lambda)_{bc} p_c \rangle = \prod_b 2\pi \delta(p_b - \text{Ad}(\lambda)_{bc} p_c). \quad (2.8)$$

Since $\text{Ad}(\lambda)_{bc}$ is diagonalized by decomposing adjoint indices into Cartan part m and ladder operator part (ij) ,

$$\begin{aligned} \langle p_b | e^{i\lambda^c G_B^c} | p_b \rangle &= (2\pi \delta(0))^{N-1} \prod_{\substack{i,j=1 \\ i \neq j}}^N 2\pi \delta([1 - z_i/z_j] p_{(ij)}) \\ &= V^{N-1} (2\pi)^{N(N-1)} \prod_{\substack{i,j=1 \\ i \neq j}}^N (1 - z_i/z_j)^{-1} \delta(p_{(ij)}), \end{aligned} \quad (2.9)$$

where V is the volume factor $V = 2\pi \delta(p = 0)$. Therefore

$$\begin{aligned} Z_B(\lambda) &= \prod_{m=1}^{N-1} \left[\int_{-\infty}^{\infty} \frac{dp_m}{2\pi} V e^{-\frac{\beta g^2}{2} p_m^2} \right] \prod_{\substack{i,j=1 \\ i \neq j}}^N (1 - z_i/z_j)^{-1} \\ &= \left(\frac{V}{\sqrt{2\pi\beta g^2}} \right)^{N-1} \prod_{\substack{i,j=1 \\ i \neq j}}^N (1 - z_i/z_j)^{-1}. \end{aligned} \quad (2.10)$$

Then Z is given by

$$\begin{aligned} Z &= \frac{1}{N!} \prod_{i=1}^{N-1} \int_0^1 dy_i \prod_{\substack{i,j=1 \\ i \neq j}}^N (z_i - z_j) \\ &\quad \times 2^{N-1} \prod_{\substack{i,j=1 \\ i \neq j}}^N (1 + z_i/z_j) \left(\frac{V}{\sqrt{2\pi\beta g^2}} \right)^{N-1} \prod_{\substack{i,j=1 \\ i \neq j}}^N (1 - z_i/z_j)^{-1} \\ &= \frac{1}{N!} \left(\frac{2V}{\sqrt{2\pi\beta g^2}} \right)^{N-1} \prod_{i=1}^{N-1} \int_0^1 dy_i \left[\prod_{\substack{i,j=1 \\ i \neq j}}^N (1 + z_i/z_j) \right]. \end{aligned} \quad (2.11)$$

Note that the factor $\prod_{i \neq j} (1 - z_i/z_j)^{-1}$ from Z_B is canceled with the measure factor from $\int d\lambda$.

To confirm the above result from different viewpoint, we will quickly explain the path integral calculation of Z (For similar calculation in a different context, see e.g. [5]). In this calculation the overall normalization of Z is somewhat ambiguous, and therefore we shall not keep track of normalization factors. In Euclidean path integral formulation Z is given by

$$Z = \frac{1}{\text{Vol}(\text{SU}(N))} \int_{\text{periodicity cond.}} DA_0 DX D\theta D\theta^\dagger \exp \left[-\frac{1}{g^2} \int_0^\beta dt \text{tr} \left[\frac{1}{2} DX DX + ig^2 \theta^\dagger D\theta - g^4 \theta^\dagger [X, \theta] \right] \right], \quad (2.12)$$

where $A_0(t)$ and $X(t)$ are periodic, and $\theta(t)$ is antiperiodic up to gauge transformation, with periodicity β . $\text{Vol}(\text{SU}(N))$ is the infinite volume of the redundancy from the gauge symmetry. By the shift $A_0 \rightarrow A_0 + g^2 X$, we can eliminate the interaction term $\theta^\dagger [X, \theta]$:

$$Z = \frac{1}{\text{Vol}(\text{SU}(N))} \int_{\text{periodicity cond.}} DA_0 DX D\theta D\theta^\dagger \exp \left[-\frac{1}{g^2} \int_0^\beta dt \text{tr} \left[\frac{1}{2} DX DX + ig^2 \theta^\dagger D\theta \right] \right]. \quad (2.13)$$

A_0 can be taken to be Δ given in the following, by taking Lorentz gauge $\partial_t A_0 = 0$:

$$\Delta = \begin{pmatrix} \alpha_1 & & & \\ & \alpha_2 & & \\ & & \ddots & \\ & & & \alpha_N \end{pmatrix}, \quad \begin{aligned} 0 \leq \alpha_1 \leq \alpha_2 \leq \cdots \leq \alpha_{N-1} < 2\pi/\beta, \\ \alpha_N = -(\alpha_1 + \alpha_2 + \cdots + \alpha_{N-1}), \\ \alpha_i : t\text{-independent}, \end{aligned} \quad (2.14)$$

General A_0 is in the form of gauge transformation of Δ : $A_0(t) = U(t)\Delta U^{-1}(t) - i\partial_t U(t)U^{-1}(t)$, and the path integral measure DA_0 is decomposed as follows:

$$DA_0 = DU \left[\prod_{i=1}^{N-1} d\alpha_i \right] \text{Det}'[D], \quad (2.15)$$

where Det' is the functional determinant for periodic functions with the zero eigenvalues removed. Although we will see that the Faddeev-Popov determinant $\text{Det}'[D]$ cancels, let us see how this is computed. By decomposing the adjoint index a into Cartan subalgebra part m and ladder operator part (ij) ,

$$\text{Det}'[D] = \prod_{m=1}^{N-1} \text{Det}'[\partial_t] \prod_{\substack{i,j=1 \\ i \neq j}}^N \text{Det}[\partial_t - i(\alpha_i - \alpha_j)]. \quad (2.16)$$

For periodic functions eigenvalues of ∂_t are given by $i\frac{2\pi}{\beta}n$ ($n \in \mathbb{Z}$) and

$$\text{Det}'[\partial_t] = \prod_{n \neq 0} i\frac{2\pi}{\beta}n = -i\beta, \quad (2.17)$$

where we used the following zeta-function regularization:

$$\prod_{n=1}^{\infty} n^{\alpha} = (2\pi)^{\alpha/2}, \quad \prod_{n=-\infty}^{\infty} a = 1. \quad (2.18)$$

Using the above and $\prod_{n=1}^{\infty} (1 - \frac{x^2}{n^2}) = \frac{\sin \pi x}{\pi x}$, $\prod_{i \neq j} \text{Det}[\partial_t - i(\alpha_i - \alpha_j)]$ can be computed similarly, and then

$$\text{Det}'[D] \sim \prod_{\substack{i,j=1 \\ i \neq j}}^N (z_i - z_j), \quad (2.19)$$

where $z_i = e^{2\pi i y_i}$, and $y_i = \frac{\beta}{2\pi} \alpha_i$. This is the measure factor for the integration over $\text{SU}(N)$ group. The path integral of DU cancels the infinite volume factor: $\int DU = \text{Vol}(\text{SU}(N))$. Since α_i appear symmetrically, the integral of α_i can be done by regarding all the α_i independent from each other, and dividing by the factor $1/N!$:

$$\frac{1}{N!} \prod_{i=1}^{N-1} \int_0^1 dy_i. \quad (2.20)$$

After the gauge fixing X and θ can be taken as periodic and antiperiodic respectively. The functional integral of them can be done as Gaussian integrals:

$$\int DX \rightarrow V^{N-1} \text{Det}'[D]^{-1}, \quad (2.21)$$

$$\int D\theta D\theta^\dagger \rightarrow \prod_{m=1}^{N-1} \widetilde{\text{Det}}[\partial_t] \prod_{\substack{i,j=1 \\ i \neq j}}^N \widetilde{\text{Det}}[\partial_t - i(\alpha_i - \alpha_j)] \sim \prod_{\substack{i,j=1 \\ i \neq j}}^N (z_i + z_j), \quad (2.22)$$

where $\widetilde{\text{Det}}$ is the functional determinant for antiperiodic functions, and the volume factor V^{N-1} comes from the integral of the zero modes of X . Then

$$\begin{aligned} Z &\sim \frac{1}{N!} \prod_{i=1}^{N-1} \int_0^1 dy_i \text{Det}'[D] \cdot V^{N-1} \text{Det}'[D]^{-1} \prod_{\substack{i,j=1 \\ i \neq j}}^N (z_i + z_j) \\ &= \frac{1}{N!} V^{N-1} \prod_{i=1}^{N-1} \int_0^1 dy_i \left[\prod_{\substack{i,j=1 \\ i \neq j}}^N (z_i + z_j) \right]. \end{aligned} \quad (2.23)$$

Thus we can reproduce the result of operator formalism calculation, with the similar cancellation of the measure factor.

To interpret the form of Z and read off information on the spectrum, let us consider mass deformation of this quantum mechanics given by the following action:

$$S = \frac{1}{g^2} \int dt \text{tr} \left[\frac{1}{2} DXDX - \frac{1}{2} g^4 M^2 X^2 + i g^2 \theta^\dagger D\theta - g^4 M \theta^\dagger \theta - g^4 \theta^\dagger [X, \theta] \right]. \quad (2.24)$$

The supersymmetry transformation is also deformed:

$$\delta X = i(\epsilon^\dagger \theta + \epsilon \theta^\dagger), \quad \delta \theta = (iMX - \Pi) \epsilon, \quad \delta A_0 = g^2 \delta X. \quad (2.25)$$

Hamiltonian and supercharge are

$$H = g^2 \text{tr} \left[\frac{1}{2} \Pi^2 + \frac{1}{2} M^2 X^2 + M \theta^\dagger \theta \right] \\ + (\text{terms proportional to 1st class constraints}), \quad (2.26)$$

$$Q = -\text{tr} [(\Pi + iMX) \theta]. \quad (2.27)$$

These satisfy the same commutation relations as those in the undeformed case. Again H acts as free Hamiltonian $h = g^2 \text{tr} \left[\frac{1}{2} \Pi^2 + \frac{1}{2} M^2 X^2 + M \theta^\dagger \theta \right]$ on the physical states. This system is easier than the undeformed case: Introducing annihilation operators $A_a = \frac{1}{\sqrt{2M}} \Pi_a - i \sqrt{\frac{M}{2}} X_a$, h is rewritten as

$$h = Mg^2 (A_a^\dagger A_a + \theta_a^\dagger \theta_a). \quad (2.28)$$

Therefore this system is equivalent to a supersymmetric harmonic oscillator system, with the physical condition $G|\text{phys}\rangle = 0$. The spectrum of states is constructed by acting $A_a^\dagger$ and $\theta_a^\dagger$ on the vacuum $|0\rangle$ defined by $A_a|0\rangle = 0$ and $\theta_a|0\rangle = 0$. For states in the following form:

$$A_{a_1}^\dagger \dots A_{a_n}^\dagger \theta_{b_1}^\dagger \dots \theta_{b_m}^\dagger |0\rangle, \quad (2.29)$$

indices $a_1, \dots, a_n$ can be symmetrized and $b_1, \dots, b_m$ antisymmetrized, and gauge invariant combinations of indices give physical states. The number of such gauge invariant states can be computed by

$$\int d\text{SU}(N) \chi(\text{Sym}_n(\text{adj.})) \chi(\text{Alt}_m(\text{adj.})). \quad (2.30)$$

Therefore the partition function of this system $Z_M = \text{Tr}_{\text{phys}}[e^{-\beta H}]$ can be computed by noting that the factor $e^{-\beta H}$ gives $q = e^{-\beta g^2 M}$ for each $A_a^\dagger$ and $\theta_a^\dagger$ on $|0\rangle$:

$$Z_M = \int d\text{SU}(N) \sum_{n=0}^{\infty} q^n \chi(\text{Sym}_n(\text{adj.})) \sum_{n=0}^{\infty} q^n \chi(\text{Alt}_n(\text{adj.})) \\ = \frac{1}{N!} \int \prod_{i=1}^{N-1} dy_i \left[\prod_{\substack{i,j=1 \\ i \neq j}}^N (1 - z_i/z_j) \right] \left(\frac{1+q}{1-q} \right)^{N-1} \prod_{\substack{i,j=1 \\ i \neq j}}^N \frac{1 + qz_i/z_j}{1 - qz_i/z_j}, \quad (2.31)$$

where we used (A.22) and (A.23). For some analysis of this expression see [6].

If we take massless limit $q \rightarrow 1$, we obtain almost the same expression of the partition function as that in the undeformed case: the factor $\prod_{i \neq j} (1 - z_i/z_j)$ is canceled, and the factor

$(1 - q)^{1-N}$ corresponds to the infinite volume factor V^{N-1} . Keeping this in mind, let us return to the structure of the spectrum in the undeformed case and try to understand the structure of Z more. If we have an energy eigenstate $|E\rangle$ with zero fermion number, we obtain new energy eigenstates by acting gauge invariant operators made of Π and $\theta^\dagger$ on $|E\rangle$, because those operators commute with the Hamiltonian[2]. Those operators can be expressed as products of single trace operators such as $\text{tr}[\Pi^n]$, $\text{tr}[\Pi^n \theta^\dagger]$, or $\text{tr}[\Pi^n \theta^\dagger \Pi^m \theta^\dagger]$. Note that at this stage we do not know if there are states which cannot be constructed in this way from a solution with zero fermion number.

We can see a correspondence of states between mass deformed and undeformed systems:

$$A_{a_1}^\dagger \dots A_{a_n}^\dagger \theta_{b_1}^\dagger \dots \theta_{b_m}^\dagger |0\rangle \longleftrightarrow \Pi_{a_1} \dots \Pi_{a_n} \theta_{b_1}^\dagger \dots \theta_{b_m}^\dagger |E\rangle. \quad (2.32)$$

Therefore number of these states in the undeformed case can be counted by setting $q = 1$ in (2.31). However, due to the factor $(1 - q)^{-1}$ we cannot take the limit $q \rightarrow 1$. This is not surprising, because there are infinitely many gauge invariant operators made of Π . This problem can be avoided as follows: Since $\text{tr}[\Pi^3]$, $\text{tr}[\Pi^4]$, $\dots$, and $\text{tr}[\Pi^N]$ are hermitian and commute with each other, h and G , these are simultaneously diagonalizable. (Note that in $\text{SU}(N)$ case $\text{tr}[\Pi^n]$ with $n > N$ can be expressed by products of those with lower n .) Therefore we consider an eigenstate $|p_2, p_3, \dots, p_N\rangle$ defined by

$$\text{tr}[\Pi^n] |p_2, p_3, \dots, p_N\rangle = p_n |p_2, p_3, \dots, p_N\rangle, \quad G^a |p_2, p_3, \dots, p_N\rangle = 0. \quad (2.33)$$

Since $\text{tr}[\Pi^n]$ do not produce new states when they act on $|p_2, p_3, \dots, p_N\rangle$, we want to remove products of $\text{tr}[\Pi^n]$ from the list of the gauge invariant operators. The list of those operators can be given by expanding the following expression:

$$\begin{aligned} & [1 + q^2 \text{tr}[\Pi^2] + (q^2 \text{tr}[\Pi^2])^2 + (q^2 \text{tr}[\Pi^2])^3 + \dots] \\ & \times [1 + q^3 \text{tr}[\Pi^3] + (q^3 \text{tr}[\Pi^3])^2 + (q^3 \text{tr}[\Pi^3])^3 + \dots] \\ & \dots \\ & \times [1 + q^N \text{tr}[\Pi^N] + (q^N \text{tr}[\Pi^N])^2 + (q^N \text{tr}[\Pi^N])^3 + \dots] \\ & = \prod_{n=2}^N \sum_{m=0}^{\infty} (q^n \text{tr}[\Pi^n])^m, \end{aligned} \quad (2.34)$$

where q is introduced to count the number of Π . If one just want to count the number of these gauge invariant operators, $\text{tr}[\Pi^n]$ in the above expression can be replaced by 1:

$$\prod_{n=2}^N \sum_{m=0}^{\infty} (q^n)^m = \prod_{n=2}^N (1 - q^n)^{-1} = (1 - q)^{1-N} \prod_{n=2}^N (1 + q + q^2 + \dots + q^{n-1})^{-1}. \quad (2.35)$$

The coefficient of q^n in the expansion of this expression gives the number of the operators made of n Π s.

Therefore the number of gauge invariant operators which act nontrivially on $|p_2, p_3, \dots, p_N\rangle$ can be counted by

$$\begin{aligned}
& \left[\int d\text{SU}(N) \sum_{n=0}^{\infty} q^n \chi(\text{Sym}_n(\text{adj.})) \sum_{n=0}^{\infty} q^n \chi(\text{Alt}_n(\text{adj.})) \right] / \\
& (1-q)^{1-N} \prod_{n=2}^N (1+q+q^2+\dots+q^{n-1})^{-1} \\
& = \frac{1}{N!} \int_0^1 \prod_{i=1}^{N-1} dy_i \left[\prod_{\substack{i,j=1 \\ i \neq j}}^N (1 - z_i/z_j) \right] (1+q)^{N-1} \prod_{n=2}^N (1+q+q^2+\dots+q^{n-1}) \\
& \times \prod_{\substack{i,j=1 \\ i \neq j}}^N \frac{1+qz_i/z_j}{1-qz_i/z_j}. \tag{2.36}
\end{aligned}$$

We can safely take the limit $q \rightarrow 1$ in this expression:

$$\int_0^1 \prod_{i=1}^{N-1} dy_i 2^{N-1} \prod_{\substack{i,j=1 \\ i \neq j}}^N (1 + z_i/z_j). \tag{2.37}$$

Note that this expression is very close to Z . (2.37) times the contribution from integrals of parameters $p_2, p_3, \dots, p_N$ gives the contribution to the partition function from states constructed by acting the gauge invariant operators on $|p_2, p_3, \dots, p_N\rangle$. At this stage we do not know how many independent solutions with the same eigenvalues of $\text{tr}[\Pi^n]$ exist in the sector of zero fermion number.

CHS solution gives a hint to the contribution from integrals of parameters $p_2, p_3, \dots, p_N$. It has $N-1$ continuous parameters $k_1, k_2, \dots, k_{N-1}$ corresponding to free particle momenta, and these appear symmetrically in the solution. Therefore the contribution to the partition function is computed by regarding k_i independent, with the factor $1/N!$:

$$\frac{1}{N!} \left[\int \frac{dk}{2\pi} \langle k | e^{-\frac{1}{2}\beta g^2 k^2} | k \rangle \right]^{N-1} = \frac{1}{N!} \left[V \int \frac{dk}{2\pi} e^{-\frac{1}{2}\beta g^2 k^2} \right]^{N-1} = \frac{1}{N!} \left[\frac{V}{\sqrt{2\pi\beta g^2}} \right]^{N-1}. \tag{2.38}$$

This times (2.37) gives exactly the same expression as Z :

$$\frac{1}{N!} \left[\frac{2V}{\sqrt{2\pi\beta g^2}} \right]^{N-1} \int_0^1 \prod_{i=1}^{N-1} dy_i \left[\prod_{\substack{i,j=1 \\ i \neq j}}^N (1 + z_i/z_j) \right]. \tag{2.39}$$

Thus we have given evidence for the following naively conjectured structure of the spectrum of states:

The whole spectrum of $D = 2$ matrix quantum mechanics is constructed by acting gauge invariant operators made of Π and at least one $\theta^\dagger$ on eigenstates of $\text{tr} [\Pi^n]$ ($n = 2, 3, \dots, N$) in the sector of zero fermion number.

and if the normalization of Z inferred from CHS solution is correct, the following conjecture follows:

In the sector of zero fermion number, CHS solution is the unique eigenfunction of $\text{tr} [\Pi^2]$, $\text{tr} [\Pi^3]$, $\dots$ and $\text{tr} [\Pi^N]$ for fixed eigenvalues.

Although we do not have proofs of this conjecture for general N , for $N = 2$ case this has been shown to be correct[1, 2]. In the next section we shall give proofs for $N = 3$ and $N = 4$ cases.

3 Analysis of eigenfunctions of $\text{tr} [\Pi^n]$ in lower N cases

In this section we give proofs of the conjecture in the previous section for $N = 3$ and $N = 4$ cases i.e. we show that CHS solutions for any N are eigenfunctions of $\text{tr} [\Pi^3]$ and $\text{tr} [\Pi^4]$, and they are the unique eigenfunctions of those operators with zero fermion number for fixed eigenvalues in $N = 3$ and $N = 4$ cases.

First we show that CHS solutions are eigenfunctions of $\text{tr} [\Pi^3]$ and $\text{tr} [\Pi^4]$. Since the expression (1.12) is regarded as the restriction of the gauge invariant solution to diagonal X , we have to know how Π acts on such restricted functions. Note that on any function $f(X)$, $\Pi_{(ij)}$ can be replaced by $G_{(ij)}$ if X is set $Y \equiv \text{diag}(x_1, \dots, x_N)$ [7].

$$\Pi_{(ij)} f(X) \Big|_{X=Y} = \frac{i}{x_i - x_j} G_{(ij)} f(X) \Big|_{X=Y}, \quad (3.1)$$

where Π_a is regarded as $-i \frac{\partial}{\partial X_a}$. Therefore, action of two or more $\Pi_{(ij)}$ s on a gauge invariant function can be reduced to that of less $\Pi_{(ij)}$ s by replacing the first $\Pi_{(ij)}$ by $G_{(ij)}$, and moving it rightward using the following commutation relation until it annihilates the function:

$$[G_{(ij)}, \Pi_{(kl)}] = i \delta_{il} \delta_{jk} (\partial_i - \partial_j) + \delta_{il} \Pi_{(jk)} - \delta_{jk} \Pi_{(il)}, \quad (3.2)$$

where $\partial_i = \frac{\partial}{\partial x_i}$ and $\Pi_{(ii)} = 0$. These can be shown from (A.3) and $w_i = (\nu^i)_m X_m$. Π_m is also replaced by ∂_i with the following relation:

$$(\nu^i)_m \Pi_m = -i(\nu^i)_m \frac{\partial}{\partial X_m} = -i \sum_j (\nu^i \cdot \nu^j) \partial_j = -i(\partial_i - P), \quad (3.3)$$

where P is the center of mass momentum operator $P = \frac{1}{N} \sum_{i=1}^N \partial_i$, and summation over m is understood. Since we set the eigenvalue of P zero and P commutes with any operator dependent only on ∂_i and the differences $x_i - x_j$, we ignore terms proportional to P in the following.

Thus we can reduce action of Π s on gauge invariant functions to ∂_i . Therefore we obtain

$$\begin{aligned} \text{tr} [\Pi^2] f(X) \Big|_{X=Y} &= \left[\sum_{i=1}^N (\nu^i)_{m_1} (\nu^i)_{m_2} \Pi_{m_1} \Pi_{m_2} + \sum'_{i,j=1}^N \Pi_{(ij)} \Pi_{(ji)} \right] f(X) \Big|_{X=Y} \\ &= - \left[\sum_{i=1}^N \partial_i \partial_i + \sum'_{i,j=1}^N \frac{2}{x_i - x_j} \partial_i \right] f(X) \Big|_{X=Y}, \end{aligned} \quad (3.4)$$

$$\begin{aligned} \text{tr} [\Pi^3] f(X) \Big|_{X=Y} &= \left[\sum_{i=1}^N (\nu^i)_{m_1} (\nu^i)_{m_2} (\nu^i)_{m_3} \Pi_{m_1} \Pi_{m_2} \Pi_{m_3} \right. \\ &\quad \left. + 3 \sum'_{i,j=1}^N (\nu^i)_m \Pi_m \Pi_{(ji)} \Pi_{(ij)} + \sum'_{i,j,k=1}^N \Pi_{(ji)} \Pi_{(kj)} \Pi_{(ik)} \right] f(X) \Big|_{X=Y} \\ &= i \left[\sum_{i=1}^N (\partial_i)^3 + 3 \sum'_{i,j=1}^N \frac{3}{x_i - x_j} (\partial_i)^2 \right. \\ &\quad \left. + 3 \sum'_{i,j,k=1}^N \frac{1}{x_j - x_i} \frac{1}{x_k - x_i} \partial_i \right] f(X) \Big|_{X=Y}, \end{aligned} \quad (3.5)$$

$$\begin{aligned} \text{tr} [\Pi^4] f(X) \Big|_{X=Y} &= \left[\sum_{i=1}^N (\nu^i)_{m_1} (\nu^i)_{m_2} (\nu^i)_{m_3} (\nu^i)_{m_4} \Pi_{m_1} \Pi_{m_2} \Pi_{m_3} \Pi_{m_4} \right. \\ &\quad + 4 \sum'_{i,j=1}^N (\nu^j)_{m_1} (\nu^j)_{m_2} \Pi_{m_1} \Pi_{m_2} \Pi_{(ji)} \Pi_{(ij)} \\ &\quad + 2 \sum'_{i,j=1}^N (\nu^j)_{m_1} (\nu^i)_{m_2} \Pi_{m_1} \Pi_{m_2} \Pi_{(ji)} \Pi_{(ij)} \\ &\quad \left. + 4 \sum'_{i,j,k=1}^N (\nu^j)_m \Pi_m \Pi_{(ji)} \Pi_{(kj)} \Pi_{(ik)} \right] f(X) \Big|_{X=Y} \end{aligned}$$

$$\begin{aligned}
& + \sum'_{i,j=1}^N \Pi_{(ji)} \Pi_{(ij)} \Pi_{(ji)} \Pi_{(ij)} + 2 \sum'_{i,j,k=1}^N \Pi_{(ji)} \Pi_{(ij)} \Pi_{(kj)} \Pi_{(jk)} \\
& + \sum'_{i,j,k,l=1}^N \Pi_{(ji)} \Pi_{(kj)} \Pi_{(lk)} \Pi_{(il)} \Big] f(X) \Big|_{X=Y} \\
& = \left[\sum_{i=1}^N (\partial_i)^4 + \sum'_{i,j=1}^N \frac{4}{x_i - x_j} (\partial_i)^3 \right. \\
& \quad + 6 \sum'_{i,j,k=1}^N \frac{1}{x_j - x_i} \frac{1}{x_k - x_i} (\partial_i)^2 \\
& \quad \left. - 4 \sum'_{i,j,k,l=1}^N \frac{1}{x_j - x_i} \frac{1}{x_k - x_i} \frac{1}{x_l - x_j} \partial_i \right] f(X) \Big|_{X=Y}, \tag{3.6}
\end{aligned}$$

where $\sum'$ indicates summations with the condition that the dummy indices take different values from each other. CHS solutions have the prefactor $\mathcal{M}$. This can be moved leftward using

$$\partial_i \frac{1}{\mathcal{M}} = \frac{1}{\mathcal{M}} \left(\partial_i + \sum_{\substack{j=1 \\ j \neq i}}^N \frac{1}{x_j - x_i} \right), \tag{3.7}$$

and with the following identities:

$$\begin{aligned}
0 &= \sum'_{i,j,k=1}^N \frac{1}{(x_j - x_i)(x_k - x_i)}, \\
0 &= \sum'_{i,j,k,l=1}^N \frac{1}{(x_j - x_i)(x_k - x_i)(x_l - x_i)}, \\
0 &= \sum'_{i,j,k,l,p=1}^N \frac{1}{(x_j - x_i)(x_k - x_i)(x_l - x_i)(x_p - x_i)}, \tag{3.8}
\end{aligned}$$

we can show

$$\text{tr} [\Pi^2] \frac{1}{\mathcal{M}} f(X) \Big|_{X=Y} = -\frac{1}{\mathcal{M}} \sum_{i=1}^N (\partial_i)^2 f(X) \Big|_{X=Y}, \tag{3.9}$$

$$\text{tr} [\Pi^3] \frac{1}{\mathcal{M}} f(X) \Big|_{X=Y} = \frac{i}{\mathcal{M}} \sum_{i=1}^N (\partial_i)^3 f(X) \Big|_{X=Y}, \tag{3.10}$$

$$\text{tr} [\Pi^4] \frac{1}{\mathcal{M}} f(X) \Big|_{X=Y} = \frac{1}{\mathcal{M}} \sum_{i=1}^N (\partial_i)^4 f(X) \Big|_{X=Y}. \tag{3.11}$$

Obviously $\sum_{\sigma} \text{sgn}(\sigma) \exp \left[i \sum_{i=1}^N k_{\sigma(i)} x_i \right]$ are eigenfunctions of $\sum_{i=1}^N (\partial_i)^3$ and $\sum_{i=1}^N (\partial_i)^4$, and therefore the above equations mean that CHS solutions are eigenfunctions of (Π^3) and (Π^4) .

Next, using the method in [3], we shall show that eigenfunctions of $\text{tr} [\Pi^n]$ with zero fermion number for fixed eigenvalues are unique in $N = 3$ and $N = 4$ cases. Gauge invariant states in the sector of zero fermion number are in the following form:

$$\psi(X) = \sum_{n_2, n_3, \dots, n_N=0}^{\infty} a_{n_2, n_3, \dots, n_N} (X^2)^{n_2} (X^3)^{n_3} \dots (X^N)^{n_N}, \quad (3.12)$$

where $(A) = \text{tr} [A]$. The following expression in $(\Pi^n)\psi$,

$$(\Pi^n)[(X^2)^{n_2} (X^3)^{n_3} \dots (X^N)^{n_N}] \quad (3.13)$$

can be rewritten as a sum of terms in the form of $(X^2)^{m_2} (X^3)^{m_3} \dots (X^N)^{m_N}$. Therefore eigenvalue equations $(\Pi^n)\psi = p_n \psi$ give recurrence relations for determining $a_{n_2, n_3, \dots, n_N}$. If these relations determine them uniquely up to the overall scaling, the eigenfunction must be unique.

For $N = 2$, the recurrence relation contains a_n and a_{n+1} , which determines a_n completely in terms of a_0 [3].

For $N = 3$, case, by explicit calculation using (A.8), (A.9) and the following equations:

$$[(\Pi^2), (X^2)^n] = 2n(3 - 2n - N^2)(X^2)^{n-1} - 4ni(X^2)^{n-1}(X\Pi), \quad (3.14)$$

$$[(\Pi^2), (X^3)^n] = 9n(n-1) \left\{ \frac{1}{N}(X^2)^2 - (X^4) \right\} (X^3)^{n-2} - 6ni(X^3)^{n-1}(X^2\Pi), \quad (3.15)$$

$$[(X\Pi), (X^3)^n] = -3ni(X^3)^n, \quad (3.16)$$

$$[(\Pi^3), (X^2)^n] = 8in(n-1)(n-2)(X^3)(X^2)^{n-3} - 6ni(X^2)^{n-1}(X\Pi^2) - 12n(n-1)(X^2)^{n-2}(X^2\Pi), \quad (3.17)$$

$$\begin{aligned} [(\Pi^3), (X^3)^n] &= i \left\{ 3n \left(\frac{4}{N} - 5N + N^3 \right) - 27n(n-1) \left(\frac{4}{N} - N \right) \right\} (X^3)^{n-1} \\ &\quad + 27in(n-1)(n-2) \left\{ \frac{2}{N^2}(X^2)^3 - \frac{3}{N}(X^2)(X^4) + (X^6) \right\} (X^3)^{n-3} \\ &\quad - 9ni(X^3)^{n-1}(X^2\Pi^2) + \frac{9n}{N}i(X^2)(X^3)^{n-1}(\Pi^2) \\ &\quad + 9n \left(\frac{4}{N} - N \right) (X^3)^{n-1}(X\Pi) - 27n(n-1)(X^3)^{n-2}(X^4\Pi) \\ &\quad + \frac{54}{N}n(n-1)(X^2)(X^3)^{n-2}(X^2\Pi), \end{aligned} \quad (3.18)$$

$$[(X\Pi^2), (X^3)^n] = \left\{ 3n \left(\frac{4}{N} - N \right) + \frac{18}{N} n(n-1) \right\} (X^2)(X^3)^{n-1} - 9n(n-1)(X^5)(X^3)^{n-2} \\ - 6ni(X^3)^{n-1}(X^3\Pi) + \frac{6n}{N} i(X^2)(X^3)^{n-1}(X\Pi), \quad (3.19)$$

$$[(X^2\Pi), (X^3)^n] = -3ni(X^4)(X^3)^{n-1} + \frac{3n}{N} i(X^2)^2(X^3)^{n-1}, \quad (3.20)$$

and the following equations which hold in $N = 3$ case,

$$(X^4) = \frac{1}{2}(X^2)^2, \quad (X^5) = \frac{5}{6}(X^2)(X^3), \quad (X^6) = \frac{1}{4}(X^2)^3 + \frac{1}{3}(X^3)^2, \quad (3.21)$$

we obtain two recurrence relations, respectively from $(\Pi^2)\psi = p_2\psi$ and $(\Pi^3)\psi = p_3\psi$:

$$p_2 a_{n,m} = -4(n+1)(n+3m+4)a_{n+1,m} - \frac{3}{2}(m+1)(m+2)a_{n-2,m+2}, \quad (3.22)$$

$$-ip_3 a_{n,m} = (m+1)\{30n+9nm+6n(n-1)+9m(m+4)+40\}a_{n,m+1} \\ - \frac{3}{4}(m+1)(m+2)(m+3)a_{n-3,m+3} \\ + 8(n+1)(n+2)(n+3)a_{n+3,m-1}, \quad (3.23)$$

where $n, m \geq 0$ and terms proportional to $a_{n',m'}$ with negative n' or m' must be omitted. From (3.22) for $n = 0$ and 1,

$$a_{1,m} = -\frac{p_2}{4(3m+4)}a_{0,m}, \quad a_{2,m} = -\frac{p_2}{8(3m+5)}a_{1,m} = \frac{(p_2)^2}{32(3m+4)(3m+5)}a_{0,m}, \quad (3.24)$$

which means that $a_{1,m}$ and $a_{2,m}$ are determined by $a_{0,m}$. Furthermore by the following relation from (3.22) for $n \geq 3$:

$$a_{n,m} = -\frac{1}{4n(n+3m+3)} \left\{ p_2 a_{n-1,m} + \frac{3}{2}(m+1)(m+2)a_{n-3,m+2} \right\}, \quad (3.25)$$

$a_{n,m}$ with $n \geq 3$ can also be determined recursively from $a_{0,m}$. Therefore there are infinitely many solutions to $(\Pi^2)\psi = p_2\psi$ and are determined by giving $a_{0,m}$ as an initial condition.

Next we shall analyze (3.23). For $n = 0$ and $m \geq 1$,

$$-ip_3 a_{0,m} = (m+1)\{9m(m+4)+40\}a_{0,m+1} + 48a_{3,m-1}. \quad (3.26)$$

Eliminating $a_{3,m-1}$ in this equation by using (3.25),

$$-ip_3 a_{0,m} = (m+1)\{9m(m+4)+40\}a_{0,m+1} - \frac{4}{3m+3} \left\{ p_2 a_{2,m-1} + \frac{3}{2}m(m+1)a_{0,m+1} \right\}, \quad (3.27)$$

and eliminating $a_{2,m-1}$ again in this equation by using (3.24),

$$0 = [(m+1)\{9m(m+4)+40\} - 2m]a_{0,m+1} + ip_3 a_{0,m} - \frac{(p_2)^3}{8(3m+1)(3m+2)(3m+3)}a_{0,m-1}. \quad (3.28)$$

This means that $a_{0,m}$ with $m \geq 2$ are determined from $a_{0,0}$ and $a_{0,1}$. Again from (3.23) for $m = 0$ and $n = 0, 1, 2$,

$$a_{0,1} = -\frac{ip_3}{40}a_{0,0}, \quad a_{1,1} = -\frac{ip_3}{70}a_{1,0}, \quad a_{2,1} = -\frac{ip_3}{112}a_{2,0}. \quad (3.29)$$

The first equation of the above means that the entire solution is determined by $a_{0,0}$ and shows that the solution for fixed p_2 and p_3 is unique. The second and third ones are consistent with (3.24). Since it seems difficult to give simple expression of the solution to (3.28), we only show some of $a_{n,m}$:

$$a_{0,2} = \frac{p_2^3 - 24p_3^2}{161280}a_{0,0}, \quad a_{0,3} = -\frac{ip_3(p_2^3 - 12p_3^2)}{35481600}a_{0,0}, \quad a_{0,4} = \frac{p_2^6 - 72p_2^3p_3^2 + 576p_3^4}{1549836288000}a_{0,0}, \quad \dots \quad (3.30)$$

This solution must coincide with CHS solution. Indeed by straightforward calculation we have checked that lower terms in the expansion of CHS solution reproduce the above $a_{n,m}$.

Thus we see that for $N = 3$ our conjecture given in the previous section is correct. Next we shall analyze $N = 4$ case. By the calculation similar to $N = 3$ case we obtain three recurrence relations:

$$\begin{aligned} p_2 a_{n,m,l} &= 2(l+1)(l+2)a_{n-3,m,l+2} + \frac{9}{4}(m+1)(m+2)a_{n-2,m+2,l} \\ &\quad - (l+1)(14m+121+29)a_{n-1,m,l+1} - \frac{4}{3}(l+1)(l+2)a_{n,m-2,l+2} \\ &\quad - 9(m+1)(m+2)a_{n,m+2,l-1} - 2(n+1)(15+2n+6m+8l)a_{n+1,m,l}, \quad (3.31) \\ -ip_3 a_{n,m,l} &= 8(n+1)(n+2)(n+3)a_{n+3,m-1,l} + 36(n+1)(n+2)(m+1)a_{n+2,m+1,l-1} \\ &\quad + 4(n+1)(l+1)(12+2l+3m+7n)a_{n+1,m-1,l+1} \\ &\quad + 9(m+1)(15+18l+4l^2+8m+4lm+m^2+7n+8ln+2mn-n^2)a_{n,m+1,l} \\ &\quad - \frac{8}{9}(l+1)(l+2)(l+3)a_{n,m-3,l+3} \\ &\quad + (l+1)(l+2)(32+28n+10m+12l)a_{n-1,m-1,l+2} \\ &\quad + 9(m+1)(l+1)(3/2+l-2n)a_{n-2,m+1,l+1} \\ &\quad + 6(l+1)(l+2)(l+3)a_{n-3,m-1,l+3} - \frac{9}{2}(m+1)(l+1)(l+2)a_{n-4,m+1,l+2}, \quad (3.32) \\ p_4 a_{n,m,l} &= \frac{27}{4}(m+1)(m+2)(l+1)(l+2)a_{n-5,m+2,l+2} \\ &\quad - (l+1)(l+2)(l+3)(37+12l+26m+8n)a_{n-4,m,l+3} \\ &\quad + \frac{81}{64}(m+1)(m+2)(m+3)(m+4)a_{n-4,m+4,l} \\ &\quad + \frac{28}{3}(l+1)(l+2)(l+3)(l+4)a_{n-3,m-2,l+4} \end{aligned}$$

$$\begin{aligned}
& -\frac{135}{8}(m+1)(m+2)(l+1)(9+4l+2m)a_{n-3,m+2,l+1} \\
& +\frac{1}{4}(l+1)(l+2)(1791+1240l+192l^2+1878m+688lm \\
& +318m^2+24n-48mn-96n^2)a_{n-2,m,l+2} \\
& -\frac{81}{8}(m+1)(m+2)(m+3)(m+4)a_{n-2,m+4,l-1} \\
& +\frac{2}{9}(l+1)(l+2)(l+3)(69+36l+50m+136n)a_{n-1,m-2,l+3} \\
& +\frac{9}{4}(m+1)(m+2)(117+222l+72l^2+50m+60lm \\
& +4m^2-51n-24ln-14mn-6n^2)a_{n-1,m+2,l} \\
& +\frac{28}{27}(l+1)(l+2)(l+3)(l+4)a_{n,m-4,l+4} \\
& +\frac{1}{2}(l+1)(1335+1384l+384l^2+32l^3+1284m+764lm+64l^2m \\
& +360m^2+84lm^2+24m^3+2102n+1768ln+320l^2n+1944mn \\
& +816lmn+336m^2n+324n^2+192ln^2+168mn^2-32n^3)a_{n,m,l+1} \\
& +\frac{81}{4}(m+1)(m+2)(m+3)(m+4)a_{n,m+4,l-2} \\
& +\frac{4}{9}(l+1)(l+2)(n+1)(135+32l+18m+78n)a_{n+1,m-2,l+2} \\
& +9(m+1)(m+2)(n+1)(45+24l+14m+6n)a_{n+1,m+2,l-1} \\
& +(n+1)(n+2)(435+480l+96l^2+396m+192lm \\
& +72m^2+116n+96ln+56mn)a_{n+2,m,l} \\
& +\frac{32}{3}(l+1)(n+1)(n+2)(n+3)a_{n+3,m-2,l+1} \\
& +16(n+1)(n+2)(n+3)(n+4)a_{n+4,m,l-1}, \tag{3.33}
\end{aligned}$$

where $n, m, l \geq 0$ and terms proportional to $a_{n',m',l'}$ with negative n' , m' or l' must be omitted. From (3.31), we obtain the following equation which determines $a_{n+1,m,l}$ with $n \geq 0$, $m \geq 0$, and $l \geq 0$, in terms of $a_{n',m',l'}$ with $n' \leq n$:

$$\begin{aligned}
a_{n+1,m,l} = & \frac{1}{2(n+1)(2n+6m+8l+15)} \left[2(l+1)(l+2)a_{n-3,m,l+2} \right. \\
& +\frac{9}{4}(m+1)(m+2)a_{n-2,m+2,l} - (l+1)(14m+12l+29)a_{n-1,m,l+1} \\
& \left. -\frac{4}{3}(l+1)(l+2)a_{n,m-2,l+2} - p_2a_{n,m,l} - 9(m+1)(m+2)a_{n,m+2,l-1} \right]. \tag{3.34}
\end{aligned}$$

Therefore we only have to determine $a_{0,m,l}$. By setting $n = 0$ and $l \geq 3$ in (3.32) and using the above to rewrite $a_{n,m,l}$ with $n \geq 1$ in terms of $a_{0,m,l}$, we obtain the following equation, which

determines $a_{0,m+5,l}$ with $m \geq 0$ and $l \geq 0$ in terms of $a_{0,m',l'}$ with $m' \leq 4$:

$$\begin{aligned}
a_{0,m+5,l} = & \frac{1}{(m+1)(m+2)(m+3)(m+4)(m+5)(37+5m+8l)} \left[\right. \\
& \frac{64}{19683} (l+4)(l+5)(l+6)(l+7)(l+8)(l+9) a_{0,m-7,l+9} \\
& + \frac{16}{2187} \frac{(l+4)(l+5)(l+6)(l+7)(39+6m+8l)}{35+6m+8l} p_2 a_{0,m-5,l+7} \\
& + \frac{4}{729} \frac{(l+4)(l+5)(41+6m+8l)}{33+6m+8l} p_2^2 a_{0,m-3,l+5} \\
& + \frac{8}{6561} (l+4)(l+5)(l+6)(119064 + 69256l + 13584l^2 + 896l^3 \\
& + 53865m + 21600lm + 2160l^2m + 9450m^2 + 1890lm^2 + 540m^3) a_{0,m-3,l+6} \\
& + \frac{1}{729} \frac{(39+6m+8l)(41+6m+8l)}{(33+6m+8l)(35+6m+8l)} p_2^3 a_{0,m-1,l+3} \\
& + \frac{2}{729} \frac{(l+4)(39+6m+8l)}{35+6m+8l} (16569 + 9730l + 1920l^2 + 128l^3 \\
& + 7497m + 2952lm + 288l^2m + 1620m^2 + 324lm^2 + 108m^3) p_2 a_{0,m-1,l+4} \\
& - \frac{i}{729} (37+6m+8l)(39+6m+8l)(41+6m+8l) p_3 a_{0,m,l+3} \\
& - \frac{1}{81} \frac{(m+1)(33+3m+8l)(41+6m+8l)}{33+6m+8l} p_2^2 a_{0,m+1,l+2} \\
& - \frac{1}{81} (m+1)(5618265 + 5767832l + 2369580l^2 + 486560l^3 + 49920l^4 + 2048l^5 \\
& + 3603906m + 2956578lm + 911180l^2m + 124800l^3m + 6400l^4m \\
& + 926103m^2 + 567682lm^2 + 116580l^2m^2 + 8000l^3m^2 + 117882m^3 \\
& + 47628lm^3 + 4860l^2m^3 + 7668m^4 + 1512lm^4 + 216m^5) a_{0,m+1,l+3} \\
& \left. - \frac{1}{9} \frac{(m+1)(m+2)(m+3)(39+6m+8l)(70+9m+16l)}{35+6m+8l} p_2 a_{0,m+3,l+1} \right]. \quad (3.35)
\end{aligned}$$

For later use we give some more equations given by setting $(n, m, l) = (0, 0, 2), (0, 0, 1), (0, 3, 0), (0, 2, 0), (0, 1, 0), (0, 0, 0)$ in (3.32):

$$-ip_3 a_{0,0,2} = \frac{9}{899} p_2^2 a_{0,1,1} + \frac{184275}{341} a_{0,1,2} + \frac{324}{319} p_2 a_{0,3,0}, \quad (3.36)$$

$$-ip_3 a_{0,1,1} = -\frac{1}{15525} p_2^3 a_{0,0,1} - \frac{2024}{1395} p_2 a_{0,0,2} + \frac{40}{2001} p_2^2 a_{0,2,0} + \frac{720720}{899} a_{0,2,1}, \quad (3.37)$$

$$-ip_3 a_{0,0,1} = \frac{3}{161} p_2^2 a_{0,1,0} + \frac{6885}{23} a_{0,1,1}, \quad (3.38)$$

$$\begin{aligned}
-ip_3 a_{0,3,0} = & -\frac{8}{27621} p_2^2 a_{0,0,2} - \frac{4240}{341} a_{0,0,3} - \frac{1}{24273} p_2^3 a_{0,2,0} \\
& - \frac{780}{899} p_2 a_{0,2,1} + \frac{555120}{341} a_{0,4,0}, \quad (3.39)
\end{aligned}$$

$$-ip_3 a_{0,2,0} = -\frac{1}{12075} p_2^3 a_{0,1,0} - \frac{570}{667} p_2 a_{0,1,1} + \frac{129789}{145} a_{0,3,0}, \quad (3.40)$$

$$-ip_3 a_{0,1,0} = -\frac{1}{4845} p_2^3 a_{0,0,0} - \frac{336}{391} p_2 a_{0,0,1} + \frac{181440}{437} a_{0,2,0}, \quad (3.41)$$

$$-ip_3 a_{0,0,0} = 135 a_{0,1,0}. \quad (3.42)$$

Henceforth we only show schematic forms of equations using the following symbol:

$$[n, m, l] \equiv \text{a term proportional to } a_{n,m,l}, \quad (3.43)$$

because explicit expressions are lengthy and it is not illuminating to show details of them. However we have computed all the explicit expressions and have confirmed that there is no singular coefficient which prevents us from solving linear equations. Setting $n = 0$ in (3.33), rewriting $a_{n,m,l}$ with $n \geq 1$ in terms of $a_{0,m,l}$, and setting $m = 0, 1, 2, 3, 4$, we obtain

$$\begin{aligned} [0, 0, l] &= [0, 0, l-1] + [0, 0, l] + [0, 0, l+1] + [0, 2, l-2] + [0, 2, l-1] \\ &\quad + [0, 4, l-3] + [0, 4, l-2] + [0, 6, l-4] + [0, 8, l-5], \end{aligned} \quad (3.44)$$

$$\begin{aligned} [0, 1, l] &= [0, 1, l-1] + [0, 1, l] + [0, 1, l+1] + [0, 3, l-2] + [0, 3, l-1] \\ &\quad + [0, 5, l-3] + [0, 5, l-2] + [0, 7, l-4] + [0, 9, l-5], \end{aligned} \quad (3.45)$$

$$\begin{aligned} [0, 2, l] &= [0, 0, l+1] + [0, 0, l+2] + [0, 2, l-1] + [0, 2, l] + [0, 2, l+1] + [0, 4, l-2] \\ &\quad + [0, 4, l-1] + [0, 6, l-3] + [0, 6, l-2] + [0, 8, l-4] + [0, 10, l-5], \end{aligned} \quad (3.46)$$

$$\begin{aligned} [0, 3, l] &= [0, 1, l+1] + [0, 1, l+2] + [0, 3, l-1] + [0, 3, l] + [0, 3, l+1] + [0, 5, l-2] \\ &\quad + [0, 5, l-1] + [0, 7, l-3] + [0, 7, l-2] + [0, 9, l-4] + [0, 11, l-5], \end{aligned} \quad (3.47)$$

$$\begin{aligned} [0, 4, l] &= [0, 0, l+3] + [0, 0, l+4] + [0, 2, l+1] + [0, 2, l+2] \\ &\quad + [0, 4, l-1] + [0, 4, l] + [0, 4, l+1] + [0, 6, l-2] + [0, 6, l-1] \\ &\quad + [0, 8, l-3] + [0, 8, l-2] + [0, 10, l-4] + [0, 12, l-5]. \end{aligned} \quad (3.48)$$

Rewriting $[0, m, l]$ with $m \geq 5$ further in terms of $[0, m', l']$ with $m' = 0, 1, 2, 3, 4$, we obtain the following: from (3.44),

$$[0, 0, 0] = [0, 0, 0] + [0, 0, 1], \quad (3.49)$$

$$[0, 0, 1] = [0, 0, 0] + [0, 0, 1] + [0, 0, 2] + [0, 2, 0], \quad (3.50)$$

$$[0, 0, 2] = [0, 0, 1] + [0, 0, 2] + [0, 0, 3] + [0, 2, 0] + [0, 2, 1] + [0, 4, 0], \quad (3.51)$$

$$[0, 0, 3] = [0, 0, 2] + [0, 0, 3] + [0, 0, 4] + [0, 2, 1] + [0, 2, 2] + [0, 4, 0] + [0, 4, 1], \quad (3.52)$$

$$\begin{aligned} [0, 0, 4] &= [0, 0, 3] + [0, 0, 4] + [0, 0, 5] + [0, 1, 3] + [0, 2, 2] + [0, 2, 3] \\ &\quad + [0, 4, 1] + [0, 4, 2], \end{aligned} \quad (3.53)$$

$$\begin{aligned}
[0, 0, l] &= [0, 0, l-1] + [0, 0, l] + [0, 0, l+1] + [0, 1, l-1] + [0, 2, l-2] \\
&\quad + [0, 2, l-1] + [0, 3, l-2] + [0, 4, l-3] + [0, 4, l-2],
\end{aligned} \tag{3.54}$$

where $l \geq 5$. From (3.45),

$$[0, 1, 0] = [0, 1, 0] + [0, 1, 1], \tag{3.55}$$

$$[0, 1, 1] = [0, 1, 0] + [0, 1, 1] + [0, 1, 2] + [0, 3, 0], \tag{3.56}$$

$$[0, 1, 2] = [0, 0, 3] + [0, 1, 1] + [0, 1, 2] + [0, 1, 3] + [0, 3, 0] + [0, 3, 1], \tag{3.57}$$

$$[0, 1, 3] = [0, 0, 3] + [0, 0, 4] + [0, 1, 2] + [0, 1, 3] + [0, 1, 4] + [0, 3, 1] + [0, 3, 2], \tag{3.58}$$

$$\begin{aligned}
[0, 1, 4] &= [0, 0, 4] + [0, 0, 5] + [0, 1, 3] + [0, 1, 4] + [0, 1, 5] + [0, 2, 3] \\
&\quad + [0, 3, 2] + [0, 3, 3],
\end{aligned} \tag{3.59}$$

$$\begin{aligned}
[0, 1, l] &= [0, 0, l] + [0, 0, l+1] + [0, 1, l] + [0, 1, l+1] + [0, 2, l-1] \\
&\quad + [0, 3, l-1] + [0, 4, l-2],
\end{aligned} \tag{3.60}$$

where $l \geq 5$. From (3.46),

$$[0, 2, 0] = [0, 0, 1] + [0, 0, 2] + [0, 2, 0] + [0, 2, 1], \tag{3.61}$$

$$[0, 2, 1] = [0, 0, 2] + [0, 0, 3] + [0, 2, 0] + [0, 2, 1] + [0, 2, 2] + [0, 4, 0], \tag{3.62}$$

$$\begin{aligned}
[0, 2, 2] &= [0, 0, 3] + [0, 0, 4] + [0, 1, 3] + [0, 2, 1] + [0, 2, 2] + [0, 2, 3] \\
&\quad + [0, 4, 0] + [0, 4, 1],
\end{aligned} \tag{3.63}$$

$$\begin{aligned}
[0, 2, 3] &= [0, 0, 3] + [0, 0, 4] + [0, 0, 5] + [0, 1, 3] + [0, 1, 4] \\
&\quad + [0, 2, 2] + [0, 2, 3] + [0, 2, 4] + [0, 4, 1] + [0, 4, 2],
\end{aligned} \tag{3.64}$$

$$\begin{aligned}
[0, 2, 4] &= [0, 0, 4] + [0, 0, 5] + [0, 0, 6] + [0, 1, 4] + [0, 1, 5] \\
&\quad + [0, 2, 3] + [0, 2, 4] + [0, 2, 5] + [0, 3, 3] + [0, 4, 2] + [0, 4, 3],
\end{aligned} \tag{3.65}$$

$$\begin{aligned}
[0, 2, l] &= [0, 0, l] + [0, 0, l+1] + [0, 0, l+2] + [0, 1, l] + [0, 2, l-1] \\
&\quad + [0, 2, l] + [0, 2, l+1] + [0, 3, l-1] + [0, 4, l-2] + [0, 4, l-1],
\end{aligned} \tag{3.66}$$

where $l \geq 5$. From (3.47),

$$[0, 3, 0] = [0, 1, 1] + [0, 1, 2] + [0, 3, 0] + [0, 3, 1], \tag{3.67}$$

$$[0, 3, 1] = [0, 0, 3] + [0, 1, 2] + [0, 1, 3] + [0, 3, 0] + [0, 3, 1] + [0, 3, 2], \tag{3.68}$$

$$\begin{aligned}
[0, 3, 2] &= [0, 0, 3] + [0, 0, 4] + [0, 1, 2] + [0, 1, 3] + [0, 1, 4] + [0, 2, 3] \\
&\quad + [0, 3, 1] + [0, 3, 2] + [0, 3, 3],
\end{aligned} \tag{3.69}$$

$$[0, 3, 3] = [0, 0, 4] + [0, 0, 5] + [0, 1, 3] + [0, 1, 4] + [0, 1, 5]$$

$$+[0, 2, 3] + [0, 2, 4] + [0, 3, 2] + [0, 3, 3] + [0, 3, 4], \quad (3.70)$$

$$\begin{aligned} [0, 3, 4] &= [0, 0, 5] + [0, 0, 6] + [0, 1, 4] + [0, 1, 5] + [0, 1, 6] \\ &\quad + [0, 2, 4] + [0, 2, 5] + [0, 3, 3] + [0, 3, 4] + [0, 3, 5] + [0, 4, 3], \end{aligned} \quad (3.71)$$

$$\begin{aligned} [0, 3, l] &= [0, 0, l+1] + [0, 0, l+2] + [0, 1, l+1] \\ &\quad + [0, 1, l+2] + [0, 2, l] + [0, 2, l+1] + [0, 3, l] \\ &\quad + [0, 3, l+1] + [0, 4, l-1], \end{aligned} \quad (3.72)$$

where $l \geq 5$. From (3.48),

$$[0, 4, 0] = [0, 0, 3] + [0, 0, 4] + [0, 2, 1] + [0, 2, 2] + [0, 4, 0] + [0, 4, 1], \quad (3.73)$$

$$\begin{aligned} [0, 4, 1] &= [0, 0, 3] + [0, 0, 4] + [0, 0, 5] + [0, 1, 3] + [0, 2, 2] + [0, 2, 3] \\ &\quad + [0, 4, 0] + [0, 4, 1] + [0, 4, 2], \end{aligned} \quad (3.74)$$

$$\begin{aligned} [0, 4, 2] &= [0, 0, 3] + [0, 0, 4] + [0, 0, 5] + [0, 0, 6] + [0, 1, 3] + [0, 1, 4] \\ &\quad + [0, 2, 2] + [0, 2, 3] + [0, 2, 4] + [0, 3, 3] + [0, 4, 1] + [0, 4, 2] + [0, 4, 3], \end{aligned} \quad (3.75)$$

$$\begin{aligned} [0, 4, 3] &= [0, 0, 4] + [0, 0, 5] + [0, 0, 6] + [0, 0, 7] + [0, 1, 4] + [0, 1, 5] + [0, 2, 3] \\ &\quad + [0, 2, 4] + [0, 2, 5] + [0, 3, 3] + [0, 3, 4] + [0, 4, 2] + [0, 4, 3] + [0, 4, 4], \end{aligned} \quad (3.76)$$

$$\begin{aligned} [0, 4, 4] &= [0, 0, 5] + [0, 0, 6] + [0, 0, 7] + [0, 0, 8] + [0, 1, 5] + [0, 1, 6] + [0, 2, 4] \\ &\quad + [0, 2, 5] + [0, 2, 6] + [0, 3, 4] + [0, 3, 5] + [0, 4, 3] + [0, 4, 4] + [0, 4, 5], \end{aligned} \quad (3.77)$$

$$\begin{aligned} [0, 4, l] &= [0, 0, l+1] + [0, 0, l+2] + [0, 0, l+3] + [0, 0, l+4] + [0, 1, l+1] \\ &\quad + [0, 1, l+2] + [0, 2, l] + [0, 2, l+1] + [0, 2, l+2] + [0, 3, l] \\ &\quad + [0, 3, l+1] + [0, 4, l-1] + [0, 4, l] + [0, 4, l+1], \end{aligned} \quad (3.78)$$

where $l \geq 5$.

From (3.49)-(3.54), we see that $a_{0,0,l+1}$ is determined by $a_{0,0,l'}$ with $l' \leq l$, $a_{0,1,l'}$ with $l' \leq l-1$, $a_{0,2,l'}$ with $l' \leq l-1$, $a_{0,3,l'}$ with $l' \leq l-2$, and $a_{0,4,l'}$ with $l' \leq l-2$.

From (3.55)-(3.59), we see that $a_{0,1,l+1}$ is determined by $a_{0,0,l'}$ with $l' \leq l+1$, $a_{0,1,l'}$ with $l' \leq l$, $a_{0,2,l'}$ with $l' \leq l-1$, $a_{0,3,l'}$ with $l' \leq l-1$, and $a_{0,4,l'}$ with $l' \leq l-2$.

From (3.61)-(3.66), we see that $a_{0,2,l+1}$ is determined by $a_{0,0,l'}$ with $l' \leq l+2$, $a_{0,1,l'}$ with $l' \leq l+1$, $a_{0,2,l'}$ with $l' \leq l$, $a_{0,3,l'}$ with $l' \leq l-1$, and $a_{0,4,l'}$ with $l' \leq l-1$.

From (3.67)-(3.72), we see that $a_{0,3,l+1}$ is determined by $a_{0,0,l'}$ with $l' \leq l+2$, $a_{0,1,l'}$ with $l' \leq l+2$, $a_{0,2,l'}$ with $l' \leq l+1$, $a_{0,3,l'}$ with $l' \leq l$, and $a_{0,4,l'}$ with $l' \leq l-1$.

We find it more useful to use the following equations obtained by eliminating $[0, 0, l + 4]$ in (3.73)-(3.78) by using (3.49)-(3.54), than to use (3.73)-(3.78):

$$[0, 4, 0] = [0, 0, 2] + [0, 0, 3] + [0, 2, 1] + [0, 2, 2] + [0, 4, 0] + [0, 4, 1], \quad (3.79)$$

$$\begin{aligned} [0, 4, 1] &= [0, 0, 3] + [0, 0, 4] + [0, 1, 3] + [0, 2, 2] + [0, 2, 3] \\ &\quad + [0, 4, 0] + [0, 4, 1] + [0, 4, 2], \end{aligned} \quad (3.80)$$

$$\begin{aligned} [0, 4, 2] &= [0, 0, 3] + [0, 0, 4] + [0, 0, 5] + [0, 1, 3] + [0, 1, 4] \\ &\quad + [0, 2, 2] + [0, 2, 3] + [0, 2, 4] + [0, 3, 3] \\ &\quad + [0, 4, 1] + [0, 4, 2] + [0, 4, 3], \end{aligned} \quad (3.81)$$

$$\begin{aligned} [0, 4, 3] &= [0, 0, 4] + [0, 0, 5] + [0, 0, 6] + [0, 1, 4] + [0, 1, 5] \\ &\quad + [0, 2, 3] + [0, 2, 4] + [0, 2, 5] + [0, 3, 3] + [0, 3, 4] \\ &\quad + [0, 4, 2] + [0, 4, 3] + [0, 4, 4], \end{aligned} \quad (3.82)$$

$$\begin{aligned} [0, 4, 4] &= [0, 0, 5] + [0, 0, 6] + [0, 0, 7] + [0, 1, 5] + [0, 1, 6] \\ &\quad + [0, 2, 4] + [0, 2, 5] + [0, 2, 6] + [0, 3, 4] + [0, 3, 5] \\ &\quad + [0, 4, 3] + [0, 4, 4] + [0, 4, 5], \end{aligned} \quad (3.83)$$

$$\begin{aligned} [0, 4, l] &= [0, 0, l + 1] + [0, 0, l + 2] + [0, 0, l + 3] + [0, 1, l + 1] + [0, 1, l + 2] \\ &\quad + [0, 2, l] + [0, 2, l + 1] + [0, 2, l + 2] + [0, 3, l + 3] + [0, 3, l + 1] \\ &\quad + [0, 4, l - 1] + [0, 4, l] + [0, 4, l + 1]. \end{aligned} \quad (3.84)$$

From these, we see that $a_{0,4,l+1}$ is determined by $a_{0,0,l'}$ with $l' \leq l + 3$, $a_{0,1,l'}$ with $l' \leq l + 2$, $a_{0,2,l'}$ with $l' \leq l + 2$, $a_{0,3,l'}$ with $l' \leq l + 1$, and $a_{0,4,l'}$ with $l' \leq l$.

In summary, $a_{0,0,l+1}$, $a_{0,1,l+1}$, $a_{0,2,l+1}$, $a_{0,3,l+1}$ and $a_{0,4,l+1}$ are determined by $a_{0,0,l'}$, $a_{0,1,l'}$, $a_{0,2,l'}$, $a_{0,3,l'}$ and $a_{0,4,l'}$ with $l' \leq l$. (These can be determined in the following order: $a_{0,0,l+1}$, $a_{0,0,l+2}$, $a_{0,1,l+1}$, $a_{0,1,l+2}$, $a_{0,2,l+1}$, $a_{0,0,l+3}$, $a_{0,3,l+1}$, $a_{0,2,l+2}$, $a_{0,4,l+1}$.) Therefore all the $a_{n,m,l}$ are determined in terms of $a_{0,0,0}$, $a_{0,1,0}$, $a_{0,2,0}$, $a_{0,3,0}$, and $a_{0,4,0}$.

Finally we determine $a_{0,1,0}$, $a_{0,2,0}$, $a_{0,3,0}$, and $a_{0,4,0}$ in terms of $a_{0,0,0}$. Explicit forms of (3.49), (3.50), (3.51) and (3.67) are

$$p_4 a_{0,0,0} = \frac{29}{68} p_2^2 a_{0,0,0} + \frac{5040}{17} a_{0,0,1}, \quad (3.85)$$

$$p_4 a_{0,0,1} = \frac{1}{101745} p_2^4 a_{0,0,0} + \frac{3159}{7820} p_2^2 a_{0,0,1} + 1496 a_{0,0,2} - \frac{4080}{437} p_2 a_{0,2,0}, \quad (3.86)$$

$$p_4 a_{0,0,2} = \frac{1}{450225} p_2^4 a_{0,0,1} + \frac{24349}{61380} p_2^2 a_{0,0,2} + \frac{45486360}{9889} a_{0,0,3}$$

$$+\frac{8}{62031}p_2^3a_{0,2,0}-\frac{9360}{899}p_2a_{0,2,1}-\frac{13909320}{9889}a_{0,4,0}, \quad (3.87)$$

$$p_4a_{0,3,0} = -\frac{4}{89001}p_2^3a_{0,1,1}+\frac{36400}{37851}p_2a_{0,1,2}+\frac{619}{1276}p_2^2a_{0,3,0}+\frac{81900}{37}a_{0,3,1}. \quad (3.88)$$

Solving these and (3.36)-(3.42), we obtain

$$a_{0,1,0} = -\frac{i}{135}p_3a_{0,0,0}, \quad (3.89)$$

$$a_{0,2,0} = \frac{-243p_2^3-1748p_3^2+684p_2p_4}{97977600}a_{0,0,0}, \quad (3.90)$$

$$a_{0,3,0} = \frac{ip_3(207p_2^3+580p_3^2-540p_2p_4)}{29099347200}a_{0,0,0}, \quad (3.91)$$

$$a_{0,4,0} = [(37503p_2^6+2834208p_2^3p_3^2+4492480p_3^4-43092p_2^4p_4-7088256p_2p_3^2p_4-688176p_2^2p_4^2+1373760p_4^3)/366065131880448000]a_{0,0,0}. \quad (3.92)$$

This shows that $a_{n,m,l}$ are determined by $a_{0,0,0}$, which gives a proof of the conjecture given in the previous section for $N = 4$ case.

4 Discussion

We have computed the thermal partition function of $SU(N)$ $D = 2$ matrix quantum mechanics, and from it we have obtained some insight into the structure of the spectrum. We have conjectured that CHS solution is the unique eigenstate of $\text{tr} [\Pi^2]$, $\text{tr} [\Pi^3]$, $\dots$, and $\text{tr} [\Pi^N]$ if we fix the eigenvalues, and have given proofs of it for $N = 3$ and $N = 4$ cases.

Our proofs of the uniqueness is too inefficient to apply to higher N cases, and it is desirable to invent more efficient method for investigating those cases. From (3.9)-(3.11), we can surmise that the following equation holds for any n :

$$\text{tr} [\Pi^n] \frac{1}{\mathcal{M}} f(X) \Big|_{X=Y} = \frac{(-i)^n}{\mathcal{M}} \sum_{i=1}^N (\partial_i)^n f(X) \Big|_{X=Y}, \quad (4.1)$$

and if this can be shown, CHS solution is proven to be an eigenfunction of $\text{tr} [\Pi^2]$, $\text{tr} [\Pi^3]$, $\dots$, and $\text{tr} [\Pi^N]$.

This system has been studied in the view of cut Fock space approach in [4, 8], which is a numerically tractable method, and some energy eigenfunctions have been constructed. Relation between it and our analysis is not immediately clear to us. It is important to make the relation clear and develop methods for numerical calculation more.

Although one may think that this system is too simplified for a model of Matrix theory, it may give hint to the structure of asymptotic plane waves of Matrix theory, which is important to consider scattering process[9].

Appendix

A $SU(N)$ group and characters

A Cartan subalgebra $\{H_m; m = 1, 2, \dots, N-1\}$ of Lie algebra $su(N)$ can be given as a set of diagonal traceless $N \times N$ matrices. For example,

$$H_m = \frac{1}{\sqrt{m(m+1)}} \begin{pmatrix} 1 & & & & \\ & \ddots & & & \\ & & 1 & & \\ & & & -m & \\ & & & & 0 \\ & & & & & \ddots \\ & & & & & & 0 \end{pmatrix}. \quad (\text{A.1})$$

These satisfy $\text{tr}(H_m H_n) = \delta_{mn}$. Then a Cartan-Weyl basis is given by

$$\{H_m, E_{ij}; m = 1, 2, \dots, N-1, i, j = 1, 2, \dots, N, i \neq j\}, \quad (\text{A.2})$$

where E_{ij} is the matrix whose only nonzero component is at i -th row and j -th column: $(E_{ij})^i_j = 1$. Commutation relations of these operators are

$$[H_m, H_n] = 0, \quad [H_m, E_{ij}] = ((\nu^i)_m - (\nu^j)_m) E_{ij}, \quad [E_{ij}, E_{kl}] = \delta_{jk} E_{il} - \delta_{il} E_{jk}, \quad (\text{A.3})$$

where

$$\nu^i = ((H_1)^i_i, (H_2)^i_i, \dots, (H_{N-1})^i_i) \quad (i = 1, 2, \dots, N), \quad (\text{A.4})$$

are weights of fundamental representation. Note that $\sum_{i=1}^N \nu^i = 0$, and the right hand side of the last equation in (A.3) can contain H_m , because E_{ii} is diagonal and can be rewritten in terms of H_m .

An ordinary hermitian basis $\{T^a; a = 1, 2, \dots, N^2 - 1\}$ satisfying $(T^a)^\dagger = T^a$ and $\text{tr}(T^a T^b) = \delta^{ab}$ is given by

$$\{T^a; a = 1, 2, \dots, N^2 - 1\} = \{H_m, E_{ij}^+, E_{ij}^-; m = 1, 2, \dots, N-1, 1 \leq i < j \leq N\}, \quad (\text{A.5})$$

where

$$E_{ij}^+ = \frac{1}{\sqrt{2}}(E_{ij} + E_{ji}), \quad E_{ij}^- = \frac{i}{\sqrt{2}}(E_{ij} - E_{ji}). \quad (\text{A.6})$$

Due to the following:

$$\sum_{i,j} (E_{ij})^k_l (E_{ji})^{k'}_{l'} = \delta^{k'}_{l'} \delta^k_l, \quad \sum_{m=1}^{N-1} (H_m)^i_j (H_m)^{i'}_{j'} = \delta^i_j \delta^{i'}_{j'} \left(\delta^{ii'} - \frac{1}{N} \right), \quad (\text{A.7})$$

T^a satisfy $\sum_a (T^a)^i_j (T^a)^{i'}_{j'} = \delta^i_{j'} \delta^{i'}_j - \frac{1}{N} \delta^i_j \delta^{i'}_{j'}$, and therefore

$$\sum_a \text{tr} (T^a A T^a B) = \text{tr} (A) \text{tr} (B) - \frac{1}{N} \text{tr} (AB), \quad (\text{A.8})$$

$$\sum_a \text{tr} (T^a A) \text{tr} (T^a B) = \text{tr} (AB) - \frac{1}{N} \text{tr} (A) \text{tr} (B). \quad (\text{A.9})$$

An element $\theta^a T^a$ of Lie algebra $su(N)$ can be expanded in various ways. We define θ^m , $\theta^{(\pm ij)}$ and $\theta^{(ij)}$ as follows:

$$\theta^a T^a = \sum_{m=1}^{N-1} \theta^m H_m + \sum_{i < j} (\theta^{(+ij)} E_{ij}^+ + \theta^{(-ij)} E_{ij}^-) \quad (\text{A.10})$$

$$= \sum_{m=1}^{N-1} \theta^m H_m + \sum_{i \neq j} \theta^{(ij)} E_{ij}. \quad (\text{A.11})$$

Adjoint representation is given by the tensor product of fundamental and antifundamental representation with the trace part subtracted. Its representation matrices are given by the structure constant f_{abc} : for a group element $e^{i\lambda^a G^a}$, where G^a are generators and λ^a are parameters, $su(N)$ representation $\text{ad}(\lambda)$ and $SU(N)$ representation $\text{Ad}(\lambda)$ are given by

$$\text{Ad}(\lambda) = e^{i\text{ad}(\lambda)}, \quad \text{ad}(\lambda)_{ab} = i f_{acb} \lambda^c. \quad (\text{A.12})$$

When $\lambda^{(ij)} = 0$, $\text{Ad}(\lambda)$ is diagonalized by decomposing the index a into m and (ij) :

$$\text{Ad}(\lambda)_{mn} = \delta_{mn}, \quad \text{Ad}(\lambda)_{(ij)(ji)} = z_j / z_i, \quad (\text{A.13})$$

where

$$z_i = e^{2\pi i y_i}, \quad 2\pi y_i = \nu^i \cdot \lambda = \sum_{m=1}^{N-1} (\nu^i)_m \lambda_m. \quad (\text{A.14})$$

Note that z_i or y_i are not independent of each other: $\prod_{i=1}^N z_i = 1$ or $\sum_{i=1}^N y_i = 0$.

The character $\chi(R)$ of a representation R is given by a trace of a group element $g = e^{i\lambda^a G^a}$ over states in R : $\chi(R) = \text{tr}_R(g)$, and we define $\chi(R^k)$ by $\chi(R^k) = \text{tr}_R(g^k)$. Since

$\text{tr}_R(hgh^{-1}) = \text{tr}_R(g)$, we can set $\lambda^{(ij)} = 0$ without loss of generality. The characters of n -rank symmetric tensor product $\text{Sym}_n(R)$, and n -rank antisymmetric tensor product $\text{Alt}_n(R)$, of a representation R can be computed by the following Frobenius formulae:

$$\chi(\text{Sym}_n(R)) = \sum_{\substack{\sum_{k=1}^n ki_k = n \\ i_k: \text{nonnegative integer}}} \prod_{k=1}^n \frac{[\chi(R^k)]^{i_k}}{i_k! \cdot k^{i_k}}, \quad (\text{A.15})$$

$$\chi(\text{Alt}_n(R)) = \sum_{\substack{\sum_{k=1}^n ki_k = n \\ i_k: \text{nonnegative integer}}} (-1)^{n + \sum_{k=1}^n i_k} \prod_{k=1}^n \frac{[\chi(R^k)]^{i_k}}{i_k! \cdot k^{i_k}}. \quad (\text{A.16})$$

The second formula gives zero when n is larger than the dimension of R . The generating functions of these characters is expressed in the following form.

$$\sum_{n=0}^{\infty} a^n \chi(\text{Sym}_n(R)) = \exp \left(\sum_{n=1}^{\infty} \frac{a^n}{n} \chi(R^n) \right), \quad (\text{A.17})$$

$$\sum_{n=0}^{\dim R} a^n \chi(\text{Alt}_n(R)) = \exp \left(- \sum_{n=1}^{\infty} \frac{(-a)^n}{n} \chi(R^n) \right). \quad (\text{A.18})$$

Often $\chi(R^n)$ is given in the form of $\chi(R^n) = \pm \sum_i b_i^n$, and in this case,

$$\begin{aligned} \sum_{n=0}^{\infty} a^n \chi(\text{Sym}_n(R)) &= \prod_i \exp \left(\pm \sum_{n=1}^{\infty} \frac{(ab_i)^n}{n} \right) = \prod_i \exp (\mp \log(1 - ab_i)) \\ &= \prod_i (1 - ab_i)^{\mp 1}, \end{aligned} \quad (\text{A.19})$$

$$\begin{aligned} \sum_{n=0}^{\dim R} a^n \chi(\text{Alt}_n(R)) &= \prod_i \exp \left(\mp \sum_{n=1}^{\infty} \frac{(-ab_i)^n}{n} \right) = \prod_i \exp (\pm \log(1 + ab_i)) \\ &= \prod_i (1 + ab_i)^{\pm 1}. \end{aligned} \quad (\text{A.20})$$

In the case of $\text{SU}(N)$ adjoint representation, the character is given by

$$\chi(\text{adj.}) = \text{tr}(\text{Ad}(\lambda)) = \sum_{i,j=1}^N \frac{z_i}{z_j} - 1, \quad (\text{A.21})$$

and

$$\sum_{n=0}^{\infty} a^n \chi(\text{Sym}_n(\text{adj.})) = (1 - a)^{-(N-1)} \prod_{\substack{i,j=1 \\ i \neq j}}^N (1 - az_i/z_j)^{-1}, \quad (\text{A.22})$$

$$\sum_{n=0}^{\dim R} a^n \chi(\text{Alt}_n(\text{adj.})) = (1+a)^{N-1} \prod_{\substack{i,j=1 \\ i \neq j}}^N (1 + a z_i / z_j). \quad (\text{A.23})$$

The square of Vandermonde determinant $\prod_{i \neq j} (z_i - z_j)$ is the measure for the following orthogonality relation for characters:

$$\begin{aligned} \int d\text{SU}(N) (\chi(R'))^* \chi(R) &= \int d\lambda (\chi(R'))^* \chi(R) \\ &\equiv \frac{1}{N!} \left(\prod_{i=1}^{N-1} \int_0^1 dy_i \right) \left[\prod_{\substack{i,j=1 \\ i \neq j}}^N (z_i - z_j) \right] (\chi(R'))^* \chi(R) \\ &= \delta_{R',R}. \end{aligned} \quad (\text{A.24})$$

References

- [1] M. Claudson and M. B. Halpern, “*Supersymmetric Ground State Wavefunctions*”, *Nucl. Phys.* **B250** (1985) 689.
- [2] S. Samuel, “*Solutions of Extended Supersymmetric Matrix Models for Arbitrary Gauge Groups*”, *Phys. Lett.* **B411** (1997) 268, [hep-th/9705167](#).
- [3] M. Trzetrzelewski, “*Large N behavior of two dimensional supersymmetric Yang-Mills quantum mechanics*”, *J.Math.Phys.* **48** (2007) 012302, [hep-th/0608147](#).
- [4] P. Korcyl, “*Exact solutions to D=2 Supersymmetric Yang-Mills Quantum Mechanics with SU(3) gauge group*”, *Acta Phys. Pol. B Proc. Suppl.* **2** (2009) 623 [arXiv:0911.2152](#) [[hep-th](#)]; P. Korcyl, “*Gauge invariant plane-wave solutions in supersymmetric Yang-Mills quantum mechanics*”, [arXiv:1008.2975](#) [[math-ph](#)]; P. Korcyl, “*Solutions of D=2 supersymmetric Yang-Mills quantum mechanics with SU(N) gauge group*”, [arXiv:1101.0591](#) [[math-ph](#)].
- [5] X. Yin, “*Matrix Models, Integrable Structures, and T-duality of Type 0 String Theory*”, *Nucl. Phys.* **B714** (2005) 137, [hep-th/0312236](#).
- [6] M. Trzetrzelewski, “*The number of gauge singlets in supersymmetric Yang-Mills quantum mechanics*”, [arXiv:0708.2946](#) [[hep-th](#)], *Phys. Rev.* **D76** (2007) 085012.

- [7] B. de Wit, M. Luscher and H. Nicolai, “*The Supermembrane Is Unstable*”, Nucl. Phys. **B320** (1989) 135.
- [8] J. Wosiek, “*Spectra of supersymmetric Yang-Mills quantum mechanics*”, NP **B644** (2002) 85, hep-th/0203116; M. Campostrini, J. Wosiek, “*Exact Witten Index in $D=2$ supersymmetric Yang-Mills quantum mechanics*”, Phys. Lett. **B550** (2002) 121, hep-th/0209140; M. Trzetrzelewski, “*The study of $SU(3)$ super Yang-Mills quantum mechanics*”, PoS LAT2005 (2005) 275, hep-th/0509197; P. Korcyl, “*Analytic calculation of Witten index in $D=2$ supersymmetric Yang-Mills quantum mechanics*”, arXiv:1101.0668 [hep-th].
- [9] J. Plefka and A. Waldron, “*On the quantum mechanics of M (atrix) theory*”, Nucl. Phys. **B512** (1998) 460. hep-th/9710104; J. Plefka, M. Serone and A. Waldron, “*The Matrix Theory S -Matrix*”, Phys. Rev. Lett. **81** (1998) 2866 hep-th/9806081.